



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Tissue transglutaminase in fibrosis - more than an ECM crosslinker

Benn, Mario C ; Weber, Willi ; Klotzsch, Enrico ; Vogel, Viola ; Pot, Simon A

Abstract: Tissue transglutaminase (TG2) is upregulated in the pathogenesis of a wide variety of chronic diseases. In this review special emphasis will be placed on fundamental mechanisms underlying the critical role of TG2 in fibroproliferative disorders. TG2 is best known for its cross-linking capacities in the extracellular space but has many critical and multifaceted roles beyond protein cross-linking, which are driven by the conformation and specific localization of the molecule. As extracellular crosslinker TG2 promotes fibrotic disease through the storage of latent TGF- β 1 in a stiffened extracellular matrix (ECM). As membrane-bound cell adhesion cofactor and signaling protein and intracellular crosslinker or G-protein, TG2 promotes fibrotic disease through cell survival and profibrotic pathway activation on a signaling, transcriptional and translational level. Similarities between the roles that TG2 plays in scar tissue and in the tumor stroma suggest that a deeper understanding of key common pathways in disease pathogenesis and progression might lead to the identification of novel treatment targets and the development of new drugs and diagnostic methods.

DOI: <https://doi.org/10.1016/j.cobme.2019.06.003>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-171497>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Benn, Mario C; Weber, Willi; Klotzsch, Enrico; Vogel, Viola; Pot, Simon A (2019). Tissue transglutaminase in fibrosis - more than an ECM crosslinker. *BioMedical Engineering OnLine*, 10:156-164.

DOI: <https://doi.org/10.1016/j.cobme.2019.06.003>

Accepted Manuscript

Tissue transglutaminase in fibrosis - more than an ECM crosslinker

Mario C. Benn, Willi Weber, Enrico Klotzsch, Viola Vogel, Simon A. Pot

PII: S2468-4511(19)30013-3

DOI: <https://doi.org/10.1016/j.cobme.2019.06.003>

Reference: COBME 149

To appear in: *Current Opinion in Biomedical Engineering*


Received Date: 18 April 2019

Revised Date: 11 June 2019

Accepted Date: 13 June 2019

Please cite this article as: M.C. Benn, W. Weber, E. Klotzsch, V. Vogel, S.A. Pot, Tissue transglutaminase in fibrosis - more than an ECM crosslinker, *Current Opinion in Biomedical Engineering*, <https://doi.org/10.1016/j.cobme.2019.06.003>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Tissue transglutaminase in fibrosis - more than an ECM crosslinker

Mario C. Benn^{#,1}, Willi Weber[@], Enrico Klotzsch^{#, @}, Viola Vogel[#], Simon A. Pot^{#, *, 1}

[#] Laboratory of Applied Mechanobiology, Department for Health Sciences and Technology, ETH Zürich, 8093 Zürich, Switzerland

[@] Institute of Biology, Experimental Biophysics/ Mechanobiology, Humboldt Universität zu Berlin, 10115 Berlin, Germany

^{*} Ophthalmology Unit, Equine Department, Vetsuisse Faculty, University of Zurich, 8057 Zurich, Switzerland

¹ equal contribution

ORCID IDs:

Mario C. Benn: 0000-0002-5345-4554

Willi Weber: 0000-0003-4110-6744

Enrico Klotzsch: 0000-0002-7577-9042

Viola Vogel: 0000-0003-2898-7671

Simon A. Pot: 0000-0002-8782-7144

Corresponding Author:

Prof. Simon A. Pot

Ophthalmology Unit, Equine Department
Vetsuisse Faculty, University of Zurich

Winterthurerstrasse 260

CH-8057 Zürich, Switzerland

Tel.: +41-44-635-9030

Fax: +41-44-635-8940

Email: spot@vetclinics.uzh.ch

Highlights of the review:

- TG2 has critical and multifaceted roles beyond ECM crosslinking
- TG2 function depends on its localization and allosterically-regulated conformation
- TG2 is upregulated in the pathogenesis of a wide variety of chronic diseases
- Many similarities exist between the roles of TG2 in the scar tissue and tumor stroma
- TG2 facilitates both TGF- β 1 storage in the ECM and TGF- β 1 activation
- TG2 is an important cell adhesion and signaling protein at the cell membrane

Keywords:

tissue transglutaminase, TG2, tTG, cross-linking, non-enzymatic, fibrosis, ECM, TGF- β 1

Abstract:

Tissue transglutaminase (TG2) is upregulated in the pathogenesis of a wide variety of chronic diseases. In this review special emphasis will be placed on fundamental mechanisms underlying the critical role of TG2 in fibroproliferative disorders. TG2 is best known for its cross-linking capacities in the extracellular space but has many critical and multifaceted roles beyond protein cross-linking, which are driven by the conformation and specific localization of the molecule.

As extracellular crosslinker TG2 promotes fibrotic disease through the storage of latent TGF- β 1 in a stiffened extracellular matrix (ECM). As membrane-bound cell adhesion cofactor and signaling protein and intracellular crosslinker or G-protein, TG2 promotes fibrotic disease through cell survival and profibrotic pathway activation on a signaling, transcriptional and translational level. Similarities between the roles that TG2 plays in scar tissue and in the tumor stroma suggest that a deeper understanding of key common pathways in disease pathogenesis and progression might lead to the identification of novel treatment targets and the development of new drugs and diagnostic methods.

TG2 - more than an ECM crosslinker

Tissue transglutaminase (TG2), also named tTG, EC 2.3.2.13, is the most studied representative of a structurally and functionally related family of proteins of which nine members have been identified in humans [1,2]. TG2 is best known for its catalytic transamidation activity, resulting in the Ca^{2+} dependent post-translational formation of covalent isopeptide bonds between glutamine and lysine residues [1]. Beyond its catalytic core, TG2 consists of an N-terminal β -sandwich and two C-terminal β -barrel domains (Figure 1, left) [3–5]. Far less understood are TG2s multiple functions at the cell membrane, in the cytoplasm and in the cell nucleus, such as adhesion, migration, growth, proliferation, survival, apoptosis, differentiation and phenotype modulation [6,7]. TG2 is an integrin and syndecan-binding adhesion co-receptor for fibronectin (Fn). Approximately 5–40% of $\beta 1$ integrins are in complex with TG2 and almost all cell-membrane bound TG2 forms 1:1 complexes with integrins [8,9], which suggests a prominent role for TG2 at the cell membrane. While the focus is often on its crosslinking capacities in the extracellular space, TG2 is also active through non-enzymatic protein–protein interactions in both the extra- and intracellular space as further highlighted below. The reader is also referred to the following comprehensive review [10].

TG2 is upregulated in the pathogenesis of a wide variety of chronic diseases

In most cases, the pathophysiological significance of TG2-induced modifications remains unclear. However, it is well documented that TG2 is involved in the pathogenesis of a wide variety of diseases, most notably neoplastic and fibroproliferative, including most malignant cancers and pulmonary, kidney and cardiac fibrosis [11–14]. TG2 expression is highly correlated with cancer cell survival, malignancy, metastasis and treatment resistance [13,15]. TG2 morphologically and functionally ‘shapes’ the tumor and scar tissue stroma through ECM cross-linking and binding of TGF- $\beta 1$ to the ECM, thus priming TGF- $\beta 1$, one of the most effective profibrotic stimuli, for release and activation [16,17]. Also, through its cell attachment and signaling mediator functions and its intracellular signaling and crosslinking functions, TG2 can promote tumor cell survival and the development of fibrotic (scar) cell phenotypes [7,8,18–23]. Major attempts are thus underway to develop potent TG2 inhibitors.

Environmental sensing through major conformational TG2 changes

The biochemical functions of TG2 largely depend on its molecular conformation (Figure 1). Various environmental factors cause allosteric conformational changes and include extracellular, intracellular and intranuclear Ca^{2+} concentrations, GDP and GTP concentrations and MMP-mediated release of membrane bound TG2 into the ECM [7,9]. When bound to GTP or GDP, TG2 adopts a closed conformation with the two C-terminal β -barrel domains folded in and blocking substrate access to the catalytic site. However, with excess Ca^{2+} levels, TG2s affinity for GDP and GTP is reduced leading to a very large change in molecular shape presenting an open molecular conformation with an accessible active catalytic cross-linking site (Figure 1b) [3,24]. Low Ca^{2+} and high GDP/GTP concentrations in the cytoplasm cause TG2 to adopt a predominantly closed conformation inside the cell [25]. The conformation that TG2 adopts at the cell membrane is largely unknown. No structure has yet been solved of TG2 complexed to any binding partner even though Fn can bind to the open and closed conformations of TG2 [5]. Closed state cytosolic TG2 can bind to heparan sulfate epitopes on

exosome membrane associated Syndecan-4 molecules which have a high binding affinity for the closed conformation of TG2. Syndecan-4-dependent translocation of TG2 from the cytoplasm to the extracellular space takes place by fusion of exosomes with the outer cell membrane [9,26–28]. This mechanism was identified in fibrotic kidney disease by Furini et al. [27]. Syndecan-4-associated TG2 delivered to the cell surface can interact with integrins and fibronectin to form membrane associated protein complexes exerting cell adhesion/receptor protein functions (Figure 2, 3) [7,29]. MMP2/9 activity can lead to Syndecan-TG2 shedding from the cell surface (Figure 3)[9]. Finally, TG2 predominantly adopts an open conformation in the extracellular space due to high Ca^{2+} and low GDP/GTP concentrations [24]. Formation of two disulfide bonds between cysteine residues under oxidizing conditions at the catalytic site renders TG2 catalytically inactive in its open conformation [30]. Active oxygen reduction mechanisms in the ECM therefore promote the catalytic ECM cross-linking activity of TG2 in the extracellular space [31]. Mechanical forces within the physiological range for cells have been shown to induce allosteric conformational and functional changes in proteins [32]. Also, TG2 is concentrated at focal adhesion sites [8,18] and the integrin and Fn-binding sites are located at opposite sides of the TG2 molecule with the catalytically active site and hinge region in between (Figure 1) [4]. It is therefore likely that cells that generate mechanical forces and pull at adhesion sites that are physically connected to the extracellular matrix can induce TG2 'opening' and catalyze disulfide bond reduction, thus inducing and stabilizing an open catalytically active conformation of TG2 [3,4].

Extra- and intracellular: Transamidation and cross-linking functions of TG2 / TGF- β 1 storage in the ECM

The catalytic activity of TG2 can affect protein conformation by generating intramolecular cross-links and can catalyze the formation of covalently linked dimers, oligomers, and polymers [6]. The transamidation activity of TG2 results in protease-resistant inter- or intramolecular isopeptide bonds which effectively cross-link ECM fibrils, stiffening the ECM and protecting the ECM from proteolytic degradation [24]. A matrix reservoir of inactive TGF- β 1 is formed via TG2-mediated cross-linking of latent TGF- β 1 binding protein (LTBP-1) to the ECM (Figure 2, 3) [33,34]. ECM-bound TGF- β 1 can be activated via mechanical release (Figure 3), which is especially effective when the LTBP-1 is bound to a stiff, deformation resistant ECM that counters cell-applied forces [17,33,35,36]. As such, TGF- β 1 activation is thought to be facilitated by TG2 since TG2 cross-links LTBP to the ECM and can turn a compliant matrix into a stiffer matrix via cross-linking of various ECM components [37]. ECM prestress sensitizes latent TGF- β 1 for activation [36], which might explain increased TGF- β 1 release from stiffened fibrotic tissues. Since increased and dysregulated expression, deposition and cross-linking of ECM cause loss of vital organ function in fibrosis, TG2 and TGF- β 1 are explored as targets for the treatment of fibrotic diseases.

Cell membrane: TG2 regulates cell adhesion and ligand affinity

TG2 binds to the fibronectin type I modules FnI_{7–9} of the collagen/gelatin binding site on the Fn molecule and interacts with cell membrane β 1, β 3 and β 5 integrins [8,38–40] and/or syndecan-4 (Figure 3) [9,28,29]. Syndecan-4 and integrin signaling is coordinated via direct complex

formation of Fn-TG2, β -integrins and syndecan-4, indirect integrin cytoplasmic tail activation via Syndecan-4 activated PKC α or downstream convergence of integrin and syndecan signaling at p190RhoGAP (Figure 3) [29,41]. TG2-Fn-complexes offer cells an alternative binding site on Fn in addition to the classical integrin-Fn binding sites (e.g. RGD loop, synergy site) and facilitate RGD independent cell adhesion [28,29,42,43].

By promoting cell adhesion to Fn via cell surface integrins and syndecan-4, TG2 likely stimulates Fn fibril assembly through cell contraction-mediated Fn stretching and unmasking of Fn self-assembly sites [38,44]. Indirectly, integrin clustering and binding of TG2 at focal adhesions might concentrate TG2 catalyzed cross-linking of the dense Fn matrix at these sites as suggested by Zemskov et al. [18].

Proteolytic degradation of surface TG2 can shift cell-ECM recognition: cells cannot bind to collagen, if TG2 is bound to certain collagen-binding integrins, such as integrin $\alpha 1\beta 1$, likely due to steric hindrance [18]. MT1-MMP and/or MMP-2 degrade integrin-associated cell surface TG2 and restore $\alpha 1\beta 1$ integrin-collagen interactions, while at the same time suppressing cell adhesion and migration on fibronectin. This shift in cell-ECM recognition via dynamic regulation of surface TG2 can affect adhesion and migration of cells on various ECM ligands [7,18].

The importance of TG2 for cell adhesion is underlined by the discovery that TG2 was 1 out of only 10 proteins that consistently appeared in seven integrin adhesome datasets analyzed by Horton et al. [45].

Cell membrane: TG2 as signaling protein

Almost all cell membrane-associated TG2 is bound to integrins [8], where TG2 dimers facilitate integrin clustering formation (Figure 3)[7,19]. Further ECM components (e.g. Fn) and cell membrane protein receptors that bind next to fibronectin's RGD-specific integrin binding site (e.g. Syndecan-4, PDGF, VEGF, FGF and EGF receptors) can be engaged by these TG2-integrin complexes to execute various cell adhesion (described above) and signaling functions (Figure 3) [7,9,21,29]. TG2-integrin clusters, TG2-Fn-integrin and TG2-Fn-integrin-syndecan-4 complexes are known to activate well reviewed integrin signaling cascades, resulting in RhoA-ROCK activation, actin filamentation and downstream release and nuclear translocation of MRTF-A and YAP/TAZ (Figure 3) [7,8,19,46,47]. Nuclear MRTF-A and YAP/TAZ both act as transcription cofactors that promote expression of profibrotic genes, including α SMA, Tenascin-C and CTGF, which can influence the tumor stroma and cancer development and promote fibrotic disease [17,48,49].

Inside the cell: Enzymatic and non-enzymatic TG2 activity

With low Ca²⁺ and high GDP/GTP concentrations in the cytoplasm, TG2 binds to and hydrolyzes GTP, assumes a closed conformation and acts as a G protein. For example, TG2 binds to c-Src and PI3-kinase, facilitating c-Src-dependent phosphorylation of PI3-kinase which promotes cell survival [22]. Due to fluctuations in Ca²⁺ levels, cytoplasmic open conformation cross-linking activity of TG2 can also be observed with various specific and unspecific target proteins. For example, TG2 drives the constitutive activation of NF- κ B through its cross-linking activity, which leads to increased TGF- β 1 expression (e.g. fibrotic diseases, cancer stroma), promotion of epithelial-mesenchymal transformation (increasing tumor malignancy) and cell survival [23,49]. Similarly, RhoA cross-linking and activation leads to increased stress fiber formation [50] and

certain translation regulatory proteins like YB-1 can be crosslinked in a state that drives increased α SMA protein translation [20]. TG2 also regulates mitochondrial function and can initiate mitochondrial-driven apoptosis by crosslinking mitochondrial proteins in its open, catalytically active state [24,51]. At the same time, TG2 cross-linking stabilizes the structure of dying cells, prevents leakage of proteolytic enzymes and protects the environment of the cell from further damage [52]. Last, high TG2 levels increase autophagy of the tumor suppressor p53, which can help tumor cells escape apoptosis [53].

Nuclear TG2 has been shown to regulate gene expression via post-translational modification of transcriptional factors and related proteins, including Sp1, hypoxia inducible factor (HIF) 1 and histones [54]. The G-protein function of cytosolic TG2 typically enhances cell survival [22], whereas the transamidation activity of TG2 can lead to cell death or survival or even fibrotic changes [20,23,49–51,55], depending on the target protein(s). Nuclear localization of TG2 is generally protective against cell death [56,57].

TG2 as major player in the development and maintenance of fibroproliferative diseases

Fibroproliferative disorders cause approximately 45% of the mortality in the developed world and appear in a wide spectrum from systemic to organ-specific fibrotic diseases. Besides increased mechanical tissue tension and the presence of certain ECM components like cellular fibronectin's ED-A domain, TGF- β 1 is one of the most effective profibrotic stimuli that drive the transformation of fibroblasts into myofibroblasts [17]. Myofibroblasts are the key players within fibroproliferative disorders [17,58]. They incorporate an excess of collagen and other fibrous proteins into the ECM while expressing strong contractile alpha smooth muscle actin (α SMA) positive stress fibers which contract the fibrotic matrix into a stiff, dysfunctional scar [17,33,48,59]. Interestingly, TG2 and TGF- β 1 reinforce each other in the progressive fibrotic and tumor stroma microenvironment [49]. As described in previous sections, TG2 facilitates the activation of TGF- β 1 in the ECM [16], the activation of profibrotic signaling cascades downstream from TG2- β -integrin-containing membrane signaling complexes [7] and the translation of α SMA [20]. TGF- β 1 on the other hand promotes TG2 transcription [60,61] leading to self-amplification of TG2 and TGF- β 1 in fibrotic tissues and explaining why both are upregulated in fibroproliferative disorders. The fact that TG2 knockout mice were protected against fibrosis [62,63] and that TG2 inhibition in mouse models of fibrosis significantly reduced the fibrotic phenotype [11,12,64] further underlines the critical role that TG2 plays in the pathogenesis and maintenance of fibrosis [65]. The relative contributions of catalytic and non-catalytic TG2 functions to healthy and diseased microenvironments still remain unknown. Inhibition of TG2s catalytic function can reduce cardiac fibrosis in vivo [11,64,66]. Wang et al. and Shinde et al. suggested further reduction of fibrosis via non-catalytic TG2-mediated mechanisms: reduced externalization of TG2 as a result of decreased interaction of intracellular TG2 with exosome membrane associated syndecan-4, decreased TGF- β 1-induced myofibroblast formation and reduced activation of fibrosis-associated genes [11,64].

Outlook: TG2 as common denominator in fibroproliferative and neoplastic diseases

The consensus in cancer research seems to mirror our descriptions of the profibrotic functions of TG2 outlined above. Extracellular TG2 in its open enzymatically active conformation promotes cancer cell malignancy through crosslinking of the tumor stroma [49,55]. Intracellular

closed conformation TG2 promotes cancer cell survival and malignancy through survival pathway and EMT activation [22,49]. Stabilizing the open state of TG2 inside the cell promotes cell death and inhibits the malignant phenotype of cancer cells, thus representing a promising new avenue in the treatment of cancer [24,25]. Similarities between scar tissue and tumor stroma [67–69] suggest that the identification of key common pathways involved in the disease pathogenesis and progression might lead to the identification of novel treatment targets and the development of new drugs and diagnostic methods.

Beyond its role as ECM crosslinker, much future research is needed to understand the many roles of TG2, including cell adhesion stabilization and its nuclear functions. Also, translational research is required as just a few TG2 inhibitors are in clinical trials and none is available for clinical use [55].

Acknowledgements: This work was supported by the ETH Zurich and by Swiss National Science Foundation Grants (Grant no. 31003A_175839/1 to VV & CR32I3_156931 to VV), the Human Frontiers Science Program (Grant no. RGY0065/2017 to EK), the Volkswagenstiftung Experiment (Grant no. 95664 to EK) and by the foundation Velux Stiftung (Project n°1227 to SP).

Declarations of interest: none

Figures + legends

All figures are to be printed in color!




Figure 1:

Functional domains, binding sites and structure of tissue transglutaminase (TG2).

Beyond its catalytic core (140 - 454), TG2 consists of an N-terminal β -sandwich motif (1 - 139) and two C-terminal β -barrel domains (1: 469-591; 2: 592-687) [3–5]. The most relevant functional binding sites will be described briefly.

- A) Recent studies identified residues K30, R116 and H134 on the N-terminal β -sandwich (yellow) as critical binding partners for the 42 kDa gelatin binding domain of Fn [5]. According to Belkin et al. TG2-integrin binding is disrupted upon deletion of C-terminal β -barrel 2 (blue), which indicates that integrin binding sites reside on that domain [4]. The catalytic core region (magenta) contains the active cross-linking site with the catalytic triad residues (C277, H335, D358). GTP (nucleotide) binding takes place on the catalytic core (K173) and β -barrel 1 (green) (R476k R478, V479, M483, R580, Y583) [70]. Heparin binding sites were identified on the catalytic core and β -barrel 2 [71]. (*) Wang et al. identified an alternative heparin binding site [9]. Ca^{2+} -binding to five of six Ca^{2+} binding sites on the catalytic core fosters allosteric catalytic site activation [72]. For information regarding further binding sites ($\alpha_1\beta$ -adrenergic receptor, PLC δ_1 , nuclear export signaling peptide) the reader is referred to the following references [70,71].
- B) X-ray crystallography demonstrated that enzymatic cross-linking activity of TG2 is only possible in the Ca^{2+} -induced open-state conformation (PDB: 2Q3Z, [3]) with exposed catalytic triad residues. Nucleotide binding induces the closed-state conformation (PDB: 1KV3) with catalytic triad residues blocked by β -barrels 1 and 2 [73]. However, since TG2 has many non-enzymatic functions, “conformational state” and “activity” must be carefully distinguished [24].

Figures are adapted and modified with permission from A [70] and B [24].




Figure 2:

TG2 functions depend on its localization, as well as its allosterically-regulated conformation as upregulated in many diseases.

Resident fibroblasts deposit and remodel the ECM in healthy tissues (top and middle left) [44], whereas α SMA expressing contractile myofibroblasts are the key players in fibrotic disease (top and middle right). Myofibroblast-dominated tissues are characterized by increased ECM fiber density and cross-linking. TG2 and TGF- β 1 are both upregulated in fibrotic tissues. The open

conformation of TG2 cross-links ECM fibrils ① [1], and binds latent TGF- β 1 via LTBP-1 to the ECM ② [1,34,74]. TG2 interacts in closed or unknown conformation with cell membrane proteins ③, resulting in downstream signaling effects [9,29,41]. Cytosolic ④ and nuclear ⑤ TG2 activate intracellular signaling pathways and downstream gene expression through enzymatic (open state TG2) and non-enzymatic (closed state TG2) mechanisms.




Figure 3:

Key pro-fibrotic roles of TG2 at the cell membrane and in the ECM.

The molecular sketches do not represent molecular dimensions nor potential conformational states. Unconventional cellular release and reuptake of TG2 are not depicted in this scheme. Figures are adapted and modified with permission from Belkin et al. [7] and Robertson et al. [75].

Membrane associated functions of TG2 (lower left section of figure):

TG2-integrin clusters, TG2-FN-Integrin and TG2-FN-integrin-syndecan-4 complexes are known to activate intracellular Src-FAK and inhibit p190RhoGAP through Src, with both pathways leading to RhoA-ROCK activation [7,8,19,29,41]. RhoA/ROCK pathway activation induces the assembly of cytoplasmic monomeric G-actin into F-actin fibers and the formation of stress fibers. Changes in actin dynamics are monitored by myocardin-related transcription factor A (MRTF-A), which binds to cytoplasmic G-actin, but dissociates from F-actin. Upon actin fiber assembly MRTF-A thus uncouples from G-actin and the released MRTF-A enters the nucleus. Alternatively, Rho-signaling and cytoskeletal stress drive the downregulation of YAP/TAZ phosphorylation and the subsequent nuclear translocation and activation of YAP/TAZ. Nuclear MRTF-A and YAP/TAZ both act as transcription cofactors that promote profibrotic gene expression [46,47]. RGD independent cell adhesion to Fn depends on Fn-TG2-syndecan-4

interaction and is mediated via cytoplasmic $\alpha 5\beta 1$ integrin and syndecan-2 activation by Syndecan-4 activated PKC α [29]. Membrane-associated TG2 interacts with growth factor receptors (such as for PDGF, VEGF, FGF and EGF) and stimulates cell survival signaling pathways (e.g. FAK, Src, Akt, etc.) by promoting GF-receptor binding and activation [7,21].

Storage of latent TGF- β 1 in the ECM and mechanoactivation of TGF- β 1 (right section of figure):

TGF- β 1 is secreted by cells in complex with its latency associated propeptide (LAP), which associates with latent TGF- β 1 binding protein (LTBP-1) [76]. LTBP-1 facilitates secretion of TGF- β 1 [77] and TG2 cross-links LTBP-1 to Fn and fibrillin ECM fibers \square [16,34]. As such, a matrix reservoir of inactive TGF- β 1 is formed which is bound to the LTBP-1 and ECM and therefore unable to bind with its high affinity cell membrane receptor to activate the downstream SMAD-mediated signaling cascade [34,74]. Latent TGF- β 1 can be activated through its release from the ECM-bound TGF β -LAP-LTBP-1 complex by proteolysis \square or integrin mediated cellular contraction \square [33–35]. The N- and C-terminal matrix binding sequences of LTBP-1 [78] and integrin binding RGD sequence of LAP [79] are critical for the activation of TGF- β 1. Cell membrane associated integrins bind to the RGD sequence on LAP \square and can expose the ECM bound LTBP-1 to cellular contractile forces. This is especially effective when the LTBP-1 is bound to a stiff, deformation resistant ECM that counters cell-applied forces [17,33,35,36]. As such, TGF- β 1 activation is thought to be facilitated by TG2 since TG2 cross-links LTBP to the ECM and can stiffen a compliant matrix via ECM cross-linking [37]. For a detailed review on TGF- β -LTBP interactions we refer the reader to a recent comprehensive review published by Rifkin et al. [74].

References

1. Lorand L, Graham RM: **Transglutaminases: crosslinking enzymes with pleiotropic functions.** *Nat Rev Mol Cell Biol* 2003, **4**:140–156.
 2. Folk JE, Chung SI: **Molecular and catalytic properties of transglutaminases.** *Adv Enzymol Relat Areas Mol Biol* 1973, **38**:109–191.
 3. Pinkas DM, Strop P, Brunger AT, Khosla C: **Transglutaminase 2 undergoes a large conformational change upon activation.** *PLoS Biol* 2007, **5**:e327.
 4. Huelsz-Prince G, Belkin AM, VanBavel E, Bakker ENTP: **Activation of extracellular transglutaminase 2 by mechanical force in the arterial wall.** *J Vasc Res* 2013, **50**:383–395.
 - *5. Cardoso I, Østerlund EC, Stamnaes J, Iversen R, Andersen JT, Jørgensen TJD, Sollid LM: **Dissecting the interaction between transglutaminase 2 and fibronectin.** *Amino Acids* 2017, **49**:489–500.
- Using biochemical and biophysical approaches, this study identified the TG2 residues K30, R116 and H134 in the N-terminal domain as critical for high affinity interaction of TG2 with Fn. Interestingly, whether TG2 adopted an open or closed state conformation did not influence Fn-binding affinity.
6. Nurminkaya MV, Belkin AM: **Cellular functions of tissue transglutaminase.** *Int Rev Cell Mol Biol* 2012, **294**:1–97.
 7. Belkin AM: **Extracellular TG2: emerging functions and regulation.** *FEBS J* 2011, **278**:4704–4716.
 8. Akimov SS, Krylov D, Fleischman LF, Belkin AM: **Tissue transglutaminase is an integrin-binding adhesion coreceptor for fibronectin.** *J Cell Biol* 2000, **148**:825–838.
 9. Wang Z, Collighan RJ, Pytel K, Rathbone DL, Li X, Griffin M: **Characterization of heparin-binding site of tissue transglutaminase: its importance in cell surface targeting, matrix deposition, and cell signaling.** *J Biol Chem* 2012, **287**:13063–13083.
 10. Kanchan K, Fuxreiter M, Fésüs L: **Physiological, pathological, and structural implications of non-enzymatic protein-protein interactions of the multifunctional human transglutaminase 2.** *Cell Mol Life Sci* 2015, **72**:3009–3035.
 11. Wang Z, Stuckey DJ, Murdoch CE, Camelliti P, Lip GYH, Griffin M: **Cardiac fibrosis can be attenuated by blocking the activity of transglutaminase 2 using a selective small-molecule inhibitor.** *Cell Death Dis* 2018, **9**:613.

12. Olsen KC, Epa AP, Kulkarni AA, Kottmann RM, McCarthy CE, Johnson GV, Thatcher TH, Phipps RP, Sime PJ: **Inhibition of transglutaminase 2, a novel target for pulmonary fibrosis, by two small electrophilic molecules.** *Am J Respir Cell Mol Biol* 2014, **50**:737–747.
13. Eckert RL: **Transglutaminase 2 takes center stage as a cancer cell survival factor and therapy target.** *Mol Carcinog* 2019, doi:10.1002/mc.22986.
- **14. Philp CJ, Siebeke I, Clements D, Miller S, Habgood A, John AE, Navaratnam V, Hubbard RB, Jenkins G, Johnson SR: **Extracellular Matrix Cross-Linking Enhances Fibroblast Growth and Protects against Matrix Proteolysis in Lung Fibrosis.** *American Journal of Respiratory Cell and Molecular Biology* 2018, **58**:594–603.

The authors used a pulmonary fibrosis cell culture (IPF fibroblasts) and an in vivo murine pulmonary fibrosis model to examine the expression of TG2 and LOX and the effects of their inhibition on fibroblast growth and ECM turnover. IPF fibroblasts strongly expressed TG2 and TG2 inhibition decreased fibroblast adhesion and proliferation and increased ECM turnover in vitro. TG2 inhibition decreased the total collagen content in the lungs of treated compared to control mice.
15. Bagatur Y, Ilter Akulke AZ, Bihorac A, Erdem M, Telci D: **Tissue transglutaminase expression is necessary for adhesion, metastatic potential and cancer stemness of renal cell carcinoma.** *Cell Adh Migr* 2018, **12**:138–151.
16. Troilo H, Steer R, Collins RF, Kielty CM, Baldock C: **Independent multimerization of Latent TGF β Binding Protein-1 stabilized by cross-linking and enhanced by heparan sulfate.** *Sci Rep* 2016, **6**:34347.
- *17. Hinz B, McCulloch CA, Coelho NM: **Mechanical regulation of myofibroblast phenoconversion and collagen contraction.** *Exp Cell Res* 2019, **379**:119–128.

This recent comprehensive review summarizes how chemical and mechanical signals induce the fibroblast-to-myofibroblast transformation and focuses on mechanobiological aspects of the cell-ECM interaction.
18. Zemskov EA, Janiak A, Hang J, Waghray A, Belkin AM: **The role of tissue transglutaminase in cell-matrix interactions.** *Front Biosci* 2006, **11**:1057–1076.
19. Janiak A, Zemskov EA, Belkin AM: **Cell surface transglutaminase promotes RhoA activation via integrin clustering and suppression of the Src-p190RhoGAP signaling pathway.** *Mol Biol Cell* 2006, **17**:1606–1619.
20. Willis WL, Hariharan S, David JJ, Strauch AR: **Transglutaminase-2 mediates calcium-regulated crosslinking of the Y-Box 1 (YB-1) translation-regulatory protein in TGF β 1-activated myofibroblasts.** *Journal of Cellular Biochemistry* 2013, **114**:2753–2769.

21. Zemskov EA, Loukinova E, Mikhailenko I, Coleman RA, Strickland DK, Belkin AM: **Regulation of platelet-derived growth factor receptor function by integrin-associated cell surface transglutaminase.** *J Biol Chem* 2009, **284**:16693–16703.
22. Boroughs LK, Antonyak MA, Cerione RA: **A novel mechanism by which tissue transglutaminase activates signaling events that promote cell survival.** *J Biol Chem* 2014, **289**:10115–10125.
23. Mann AP, Verma A, Sethi G, Manavathi B, Wang H, Fok JY, Kunnumakkara AB, Kumar R, Aggarwal BB, Mehta K: **Overexpression of tissue transglutaminase leads to constitutive activation of nuclear factor-kappaB in cancer cells: delineation of a novel pathway.** *Cancer Res* 2006, **66**:8788–8795.
- **24. Katt WP, Antonyak MA, Cerione RA: **Opening up about Tissue Transglutaminase: When Conformation Matters More than Enzymatic Activity.** *Med One* 2018, **3**.

A thorough review of the different conformational states of TG2, the binding partners associated with TG2's different conformational states and the implications in health and disease. The authors elegantly state that the conformation and activity of TG2 must be considered individually. The focus of this review is on novel therapies and cancer.
25. Katt WP, Blobel NJ, Komarova S, Antonyak MA, Nakano I, Cerione RA: **A small molecule regulator of tissue transglutaminase conformation inhibits the malignant phenotype of cancer cells.** *Oncotarget* 2018, **9**:34379–34397.
26. Furini G, Verderio EAM: **Spotlight on the Transglutaminase 2-Heparan Sulfate Interaction.** *Med Sci (Basel)* 2019, **7**.
27. Furini G, Schroeder N, Huang L, Boocock D, Scarpellini A, Coveney C, Tonoli E, Ramaswamy R, Ball G, Verderio C, et al.: **Proteomic Profiling Reveals the Transglutaminase-2 Externalization Pathway in Kidneys after Unilateral Ureteric Obstruction.** *J Am Soc Nephrol* 2018, **29**:880–905.
28. Lortat-Jacob H, Burhan I, Scarpellini A, Thomas A, Imbert A, Vivès RR, Johnson T, Gutierrez A, Verderio EAM: **Transglutaminase-2 interaction with heparin: identification of a heparin binding site that regulates cell adhesion to fibronectin-transglutaminase-2 matrix.** *J Biol Chem* 2012, **287**:18005–18017.
29. Wang Z, Collighan RJ, Gross SR, Danen EHJ, Orend G, Telci D, Griffin M: **RGD-independent cell adhesion via a tissue transglutaminase-fibronectin matrix promotes fibronectin fibril deposition and requires syndecan-4/2 $\alpha 5\beta 1$ integrin co-signaling.** *J Biol Chem* 2010, **285**:40212–40229.
30. Stamnaes J, Pinkas DM, Fleckenstein B, Khosla C, Sollid LM: **Redox Regulation of Transglutaminase 2 Activity.** *Journal of Biological Chemistry* 2010, **285**:25402–25409.

31. Plugis NM, Palanski BA, Weng C-H, Albertelli M, Khosla C: **Thioredoxin-1 Selectively Activates Transglutaminase 2 in the Extracellular Matrix of the Small Intestine: IMPLICATIONS FOR CELIAC DISEASE.** *J Biol Chem* 2017, **292**:2000–2008.
32. Schoen I, Pruitt BL, Vogel V: **The Yin-Yang of Rigidity Sensing: How Forces and Mechanical Properties Regulate the Cellular Response to Materials.** *Annual Review of Materials Research* 2013, **43**:589–618.
- *33. Klingberg F, Chau G, Walraven M, Boo S, Koehler A, Chow ML, Olsen AL, Im M, Lodyga M, Wells RG, et al.: **The fibronectin ED-A domain enhances recruitment of latent TGF- β -binding protein-1 to the fibroblast matrix.** *J Cell Sci* 2018, **131**.

Stiffness-tuneable substrates for fibroblast cultures were used to demonstrate that high substrate stiffness promotes the co-expression of ED-A fibronectin and LTBP-1. By binding to LTBP-1, the ED-A Fn splice variant thus enhanced the storage of latent TGF- β 1 in the ECM.
34. Nunes I, Gleizes PE, Metz CN, Rifkin DB: **Latent transforming growth factor-beta binding protein domains involved in activation and transglutaminase-dependent cross-linking of latent transforming growth factor-beta.** *J Cell Biol* 1997, **136**:1151–1163.
35. Buscemi L, Ramonet D, Klingberg F, Formey A, Smith-Clerc J, Meister J-J, Hinz B: **The single-molecule mechanics of the latent TGF- β 1 complex.** *Curr Biol* 2011, **21**:2046–2054.
36. Klingberg F, Chow ML, Koehler A, Boo S, Buscemi L, Quinn TM, Costell M, Alman BA, Genot E, Hinz B: **Prestress in the extracellular matrix sensitizes latent TGF- β 1 for activation.** *J Cell Biol* 2014, **207**:283–297.
37. Greenberg CS, Birckbichler PJ, Rice RH: **Transglutaminases: multifunctional cross-linking enzymes that stabilize tissues.** *FASEB J* 1991, **5**:3071–3077.
38. Akimov SS, Belkin AM: **Cell-surface transglutaminase promotes fibronectin assembly via interaction with the gelatin-binding domain of fibronectin: a role in TGFbeta-dependent matrix deposition.** *J Cell Sci* 2001, **114**:2989–3000.
39. Mangala LS, Fok JY, Zorrilla-Calancha IR, Verma A, Mehta K: **Tissue transglutaminase expression promotes cell attachment, invasion and survival in breast cancer cells.** *Oncogene* 2007, **26**:2459–2470.
40. Soluri MF, Boccafroschi F, Cotella D, Moro L, Forestieri G, Autiero I, Cavallo L, Oliva R, Griffin M, Wang Z, et al.: **Mapping the minimum domain of the fibronectin binding site on transglutaminase 2 (TG2) and its importance in mediating signaling, adhesion, and migration in TG2-expressing cells.** *FASEB J* 2019, **33**:2327–2342.
41. Bass MD, Morgan MR, Roach KA, Settleman J, Goryachev AB, Humphries MJ: **p190RhoGAP is the convergence point of adhesion signals from alpha 5 beta 1 integrin and syndecan-4.** *J Cell Biol* 2008, **181**:1013–1026.

42. Verderio EAM, Telci D, Okoye A, Melino G, Griffin M: **A Novel RGD-independent Cell Adhesion Pathway Mediated by Fibronectin-bound Tissue Transglutaminase Rescues Cells from Anoikis.** *Journal of Biological Chemistry* 2003, **278**:42604–42614.
43. Telci D, Wang Z, Li X, Verderio EAM, Humphries MJ, Baccarini M, Basaga H, Griffin M: **Fibronectin-tissue transglutaminase matrix rescues RGD-impaired cell adhesion through syndecan-4 and beta1 integrin co-signaling.** *J Biol Chem* 2008, **283**:20937–20947.
44. Vogel V: **Unraveling the Mechanobiology of Extracellular Matrix.** *Annu Rev Physiol* 2018, **80**:353–387.
45. Horton ER, Byron A, Askari JA, Ng DHJ, Millon-Frémillon A, Robertson J, Koper EJ, Paul NR, Warwood S, Knight D, et al.: **Definition of a consensus integrin adhesome and its dynamics during adhesion complex assembly and disassembly.** *Nat Cell Biol* 2015, **17**:1577–1587.
46. Asparuhova MB, Gelman L, Chiquet M: **Role of the actin cytoskeleton in tuning cellular responses to external mechanical stress.** *Scand J Med Sci Sports* 2009, **19**:490–499.
47. Panciera T, Azzolin L, Cordenonsi M, Piccolo S: **Mechanobiology of YAP and TAZ in physiology and disease.** *Nat Rev Mol Cell Biol* 2017, **18**:758–770.
48. Kennedy L, Shi-Wen X, Carter DE, Abraham DJ, Leask A: **Fibroblast adhesion results in the induction of a matrix remodeling gene expression program.** *Matrix Biol* 2008, **27**:274–281.
49. Agnihotri N, Mehta K: **Transglutaminase-2: evolution from pedestrian protein to a promising therapeutic target.** *Amino Acids* 2017, **49**:425–439.
50. Singh US, Kunar MT, Kao YL, Baker KM: **Role of transglutaminase II in retinoic acid-induced activation of RhoA-associated kinase-2.** *EMBO J* 2001, **20**:2413–2423.
51. Pavlyukov MS, Antipova NV, Balashova MV, Shakhparonov MI: **Detection of Transglutaminase 2 conformational changes in living cell.** *Biochem Biophys Res Commun* 2012, **421**:773–779.
52. Park D, Choi SS, Ha K-S: **Transglutaminase 2: a multi-functional protein in multiple subcellular compartments.** *Amino Acids* 2010, **39**:619–631.
53. Kang JH, Lee J-S, Hong D, Lee S-H, Kim N, Lee W-K, Sung T-W, Gong Y-D, Kim S-Y: **Renal cell carcinoma escapes death by p53 depletion through transglutaminase 2-chaperoned autophagy.** *Cell Death Dis* 2016, **7**:e2163.
54. Kuo T-F, Tatsukawa H, Kojima S: **New insights into the functions and localization of nuclear transglutaminase 2.** *FEBS J* 2011, **278**:4756–4767.
55. Katt WP, Antonyak MA, Cerione RA: **The diamond anniversary of tissue transglutaminase: a protein of many talents.** *Drug Discov Today* 2018, **23**:575–591.

56. Milakovic T, Tucholski J, McCoy E, Johnson GVW: **Intracellular localization and activity state of tissue transglutaminase differentially impacts cell death.** *J Biol Chem* 2004, **279**:8715–8722.
57. Gundemir S, Johnson GVW: **Intracellular localization and conformational state of transglutaminase 2: implications for cell death.** *PLoS One* 2009, **4**:e6123.
58. Van De Water L, Varney S, Tomasek JJ: **Mechanoregulation of the Myofibroblast in Wound Contraction, Scarring, and Fibrosis: Opportunities for New Therapeutic Intervention.** *Adv Wound Care* 2013, **2**:122–141.
59. Goffin JM, Pittet P, Csucs G, Lussi JW, Meister J-J, Hinz B: **Focal adhesion size controls tension-dependent recruitment of alpha-smooth muscle actin to stress fibers.** *J Cell Biol* 2006, **172**:259–268.
60. Ritter SJ, Davies PJ: **Identification of a transforming growth factor-beta1/bone morphogenetic protein 4 (TGF-beta1/BMP4) response element within the mouse tissue transglutaminase gene promoter.** *J Biol Chem* 1998, **273**:12798–12806.
61. Sándor K, Daniel B, Kiss B, Kovács F, Szondy Z: **Transcriptional control of transglutaminase 2 expression in mouse apoptotic thymocytes.** *Biochim Biophys Acta* 2016, **1859**:964–974.
62. Shweke N, Boulous N, Jouanneau C, Vandermeersch S, Melino G, Dussault J-C, Chatziantoniou C, Ronco P, Boffa J-J: **Tissue Transglutaminase Contributes to Interstitial Renal Fibrosis by Favoring Accumulation of Fibrillar Collagen through TGF- β Activation and Cell Infiltration.** *The American Journal of Pathology* 2008, **173**:631–642.
63. Olsen KC, Sapinoro RE, Kottmann RM, Kulkarni AA, Iismaa SE, Johnson GVW, Thatcher TH, Phipps RP, Sime PJ: **Transglutaminase 2 and its role in pulmonary fibrosis.** *Am J Respir Crit Care Med* 2011, **184**:699–707.
64. Shinde AV, Su Y, Palanski BA, Fujikura K, Garcia MJ, Frangogiannis NG: **Pharmacologic inhibition of the enzymatic effects of tissue transglutaminase reduces cardiac fibrosis and attenuates cardiomyocyte hypertrophy following pressure overload.** *J Mol Cell Cardiol* 2018, **117**:36–48.
65. Szondy Z, Korponay-Szabó I, Király R, Sarang Z, Tsay GJ: **Transglutaminase 2 in human diseases.** *Biomedicine* 2017, **7**:15.
66. Shinde AV, Frangogiannis NG: **Tissue transglutaminase in the pathogenesis of heart failure.** *Cell Death & Differentiation* 2018, **25**:453–456.
67. Rybinski B, Franco-Barraza J, Cukierman E: **The wound healing, chronic fibrosis, and cancer progression triad.** *Physiol Genomics* 2014, **46**:223–244.

68. Cox TR, Erler JT: **Molecular pathways: connecting fibrosis and solid tumor metastasis.** *Clin Cancer Res* 2014, **20**:3637–3643.
69. Foster DS, Jones RE, Ransom RC, Longaker MT, Norton JA: **The evolving relationship of wound healing and tumor stroma.** *JCI Insight* 2018, **3**.
70. Lai T-S, Greenberg CS: **TGM2 and implications for human disease: role of alternative splicing.** *Front Biosci* 2013, **18**:504–519.
71. Scarpellini A, Germack R, Lortat-Jacob H, Muramatsu T, Billett E, Johnson T, Verderio EAM: **Heparan Sulfate Proteoglycans Are Receptors for the Cell-surface Trafficking and Biological Activity of Transglutaminase-2.** *Journal of Biological Chemistry* 2009, **284**:18411–18423.
72. Casadio R, Polverini E, Mariani P, Spinozzi F, Carsughi F, Fontana A, Polverino de Laureto P, Matteucci G, Bergamini CM: **The structural basis for the regulation of tissue transglutaminase by calcium ions.** *Eur J Biochem* 1999, **262**:672–679.
73. Begg GE, Carrington L, Stokes PH, Matthews JM, Wouters MA, Husain A, Lorand L, Iismaa SE, Graham RM: **Mechanism of allosteric regulation of transglutaminase 2 by GTP.** *Proc Natl Acad Sci U S A* 2006, **103**:19683–19688.
- *74. Rifkin DB, Rifkin WJ, Zilberberg L: **LTBPs in biology and medicine: LTBP diseases.** *Matrix Biol* 2018, **71-72**:90–99.

This excellent review summarizes the current knowledge state regarding LTBP and LTBP related disease. LTBP is a crucial mediator of TGF- β activity and a key substrate of TG2.
75. Robertson IB, Rifkin DB: **Unchaining the beast; insights from structural and evolutionary studies on TGF β secretion, sequestration, and activation.** *Cytokine Growth Factor Rev* 2013, **24**:355–372.
76. Öklü R, Hesketh R: **The latent transforming growth factor β binding protein (LTBP) family.** *Biochemical Journal* 2000, **352**:601–610.
77. Miyazono K, Olofsson A, Colosetti P, Heldin CH: **A role of the latent TGF-beta 1-binding protein in the assembly and secretion of TGF-beta 1.** *The EMBO Journal* 1991, **10**:1091–1101.
78. Mazzieri R, Jurukovski V, Obata H, Sung J, Platt A, Annes E, Karaman-Jurukovska N, Gleizes P-E, Rifkin DB: **Expression of truncated latent TGF-beta-binding protein modulates TGF-beta signaling.** *J Cell Sci* 2005, **118**:2177–2187.
79. Yang Z, Mu Z, Dabovic B, Jurukovski V, Yu D, Sung J, Xiong X, Munger JS: **Absence of integrin-mediated TGFbeta1 activation in vivo recapitulates the phenotype of TGFbeta1-null mice.** *J Cell Biol* 2007, **176**:787–793.

AUTHOR DECLARATION TEMPLATE

We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

We confirm that the manuscript has been read and approved by all named authors and that there are no other persons who satisfied the criteria for authorship but are not listed. We further confirm that the order of authors listed in the manuscript has been approved by all of us.

We confirm that we have given due consideration to the protection of intellectual property associated with this work and that there are no impediments to publication, including the timing of publication, with respect to intellectual property. In so doing we confirm that we have followed the regulations of our institutions concerning intellectual property.

We understand that the Corresponding Author is the sole contact for the Editorial process (including Editorial Manager and direct communications with the office). He/she is responsible for communicating with the other authors about progress, submissions of revisions and final approval of proofs. We confirm that we have provided a current, correct email address which is accessible by the Corresponding Author and which has been configured to accept email from

spot@vetclinics.uzh.ch

Signed by all authors as follows:

14.04.2019 Enrico Klotzsch



14.04.2019 Simon Pot



14.04.2019 Mario Benn



14.04.2019 Willi Weber



16.04.2019 Viola Vogel

